arXiv:math/0412326v1 [math.RA] 16 Dec 2004

Effective computation of $\mathrm{Tor}_k(M,N)$

Socorro García Román [*]   Manuel García Román

February 1, 2008

Abstract

An effective method to compute a presentation of $\mathrm{Tor}_k(M,N)$ for modules on a not necessarily commutative algebra is proposed.

Introduction

Unlike $\mathrm{Ext}_k(M,N)$ (see [4] for an algorithm when the base ring R is a commutative algebra and [1] when R is a PBW algebra), there are not known effective methods to compute $\mathrm{Tor}_k(M,N)$ for a pair of R-modules M and N.

In this note, we propose an algorithm to compute a presentation of $\mathrm{Tor}_k(M,N)$ when R is a Poincaré-Birkhoff-Witt algebra (PBW algebra for short; see [1, 2, et.al.] for a definition and examples, including the Weyl algebras, universal enveloping algebras of Lie algebras, the quantum plane, algebras of quantum matrices and other iterated Ore extensions, etc.). Since in general $\mathrm{Tor}_k(M,N)$ is just an abelian group when M and N are left R-modules, we ask for a two-sided structure on M. When M is an R-bimodule then $\mathrm{Tor}_k(M,N)$ is a left R-module. We show that if, in addition, M is finitely generated and N is a finitely generated left R-module, then effective techniques involving Gröbner bases may be used to compute $\mathrm{Tor}_k(M,N)$ for any $k \geq 0$.

Besides the standard algorithm to compute (left) syzygies, the ingredients of our method, are

- an algorithm for computing a free resolution of the left R-module N given a finite system of generators of N. Such an algorithm may be found in [2];

---

[*] Supported by DGUI, Consejería de Educación - Gobierno de Canarias

- a finite presentation of the R-bimodule M. Using the syzygy bimodule described in [3], we show below a method for computing such a presentation when M is a centralizing R-bimodule and a finite system of two-sided generators of M is given;
- theorem 2.2;

Our algorithm, described in detail in 2.3, follows the lines of [1].

1 Preliminaries

This section is devoted to fixing notation and describing some isomorphisms which will be used later.

Let M be an R-bimodule.

For all $s \geq 1$, the map $\alpha : M^s \longrightarrow M \otimes_R R^s$; $\alpha(f_1, \ldots, f_s) = \sum_{i=1}^{s} f_i \otimes e_i$, where $\{e_1, .., e_s\}$ is the canonical basis of R^s, is an isomorphism of left R-modules.

Indeed, α is the composition of the isomorphisms

$$\begin{array}{ccccc} M^s & \longrightarrow & (M \otimes_R R)^s & \longrightarrow & M \otimes_R R^s \\ (f_1, \ldots, f_s) & \mapsto & (f_1 \otimes 1, \ldots, f_s \otimes 1) & \mapsto & \sum_{i=1}^{s} f_i \otimes e_i. \end{array} \tag{1}$$

On the other hand, if A is a subbimodule of R^m and B is a left submodule of R^s, then

$$\begin{array}{rcl} \beta : (R^m \otimes_R R^s)/T & \longrightarrow & (R^m/A) \otimes_R (R^s/B) \\ (f \otimes g) + T & \mapsto & (f + A) \otimes (g + B), \end{array} \tag{2}$$

where $T = R^m \otimes B \ + \ A \otimes_R R^s$, is an isomorphism of left R-modules.

Furthermore, if $\{a_1, \ldots, a_r\}$ is a generator system of A such that $A = Ra_1 + \cdots + Ra_r = a_1R + \cdots + a_rR$ and $\{b_1, \ldots, b_t\}$ is a generator system of B as a left R-module, then

$$\{a_i \otimes b_j \ / \ 1 \leq i \leq r, \ 1 \leq j \leq t\}$$

is a generator system of $A \otimes_R B$ as a left R-module, since for all $a =$

$\sum_{l=l}^{r} p_l a_l \in A$ and $b = \sum_{j=1}^{t} p'_j b_j \in B$ with $p_l, p'_j \in R$,

$$\begin{aligned} a \otimes b &= \sum_{j,l} p_l a_l \otimes p'_j b_j \\ &= \sum_{j,l} p_l (a_l p'_j) \otimes b_j \\ &= \sum_{j,l} p_l (\sum_{i=1}^{r} p_i^{jl} a_i) \otimes b_j \\ &= \sum_{i,j,l} p_l p^{jl} a_i \otimes b_j . \end{aligned}$$

¿From here on, let $0 \to L \to R^m \overset{p_M}{\to} M \to 0$ be a finite presentation of the R-bimodule M, and let $H = \{h_1, \ldots, h_r\} \subseteq R^m$ be a two-sided generator system of L in such a way that $L = RH = HR$ (e.g., H is a two-sided Gröbner basis of L).

Remark 1.1. If M is a centralizing bimodule, such a presentation may be computed as follows. Recall that the R-bimodule M is said to be centralizing if M is generated as a left R-module (or equivalently, as a right R-module) by its centralizer

$$\mathrm{Cen}_R M = \{m \in M;\, rm = mr, \forall\, r \in R\}.$$

Let R be a PBW algebra and M an R-bimodule. In [3] we define the syzygy bimodule of $\{m_1, \ldots, m_s\} \subseteq M$ as the kernel of the morphism of left R^{env}-modules $(R^{\mathrm{env}})^s \longrightarrow M; \quad e_i \longmapsto m_i$, where R^{env} is the enveloping algebra $R \otimes R^{\mathrm{op}}$. We also provide an algorithm to compute the syzygy bimodule of $\{m_1, \ldots, m_s\}$.

If the R-bimodule M is centralizing, say generated by $\{m_1, \ldots, m_s\} \in \mathrm{Cen}_R M$, then the syzygy bimodule is $(\mathfrak{m}^s)^{-1} L$, where $\mathfrak{m}^s : (R^{\mathrm{env}})^s \longrightarrow R^s$ is given by $\mathfrak{m}(f \otimes g) = fg$ and L is the kernel of the epimorphism $p_M : R^s \longrightarrow M$. In that case, if G is a (left) Gröbner basis of the syzygy bimodule, then $H = \mathfrak{m}^s(G) \setminus \{0\}$ is a two-sided Gröbner basis of L (cf. [3, thm. 7]).

If N is a left R-module and $0 \to B \to R^s \to N \to 0$ is a presentation of N, then the map

$$\begin{array}{rrcl} \gamma : & R^{ms}/\alpha^{-1}(T) & \longrightarrow & M \otimes_R N \\ & e_{ij} + \alpha^{-1}(T) & \longmapsto & p_M\,(e'_j) \otimes p_N(e_i) \end{array} \tag{3}$$

is an isomorphism of left R-modules, where $T = R^m \otimes_R B \,+\, L \otimes_R R^s$, and $\{e_1, \ldots, e_s\}$, resp. $\{e'_1, \ldots, e'_m\}$, resp. $\{e_{11}, \ldots, e_{1m}, \ldots, e_{s1}, \ldots, e_{sm}\}$ are the canonical bases of R^s, resp. R^m, resp. $R^{ms} = (R^m)^s$, being $e_{ij} = (0_{R^m}, \ldots, \overset{i}{e'_j}, \ldots, 0_{R^m})$.

Indeed, γ is the composition of isomorphisms

$$R^{ms}/\alpha^{-1}(T) \xrightarrow{\bar{\alpha}} (R^m \otimes_R R^s)/T \xrightarrow{\beta} (R^m/L) \otimes_R (R^s/B) \overset{\bar{p}_M \otimes \bar{p}_N}{\longrightarrow} M \otimes_R N$$

where $\bar{\alpha}$, resp. $\bar{p_M}$, resp. $\bar{p_N}$ is obtained by factoring $\alpha : R^{ms} \longrightarrow (R^m \otimes_R R^s)/T$, resp. $p_M : R^m \longrightarrow M$, resp. $p_N : R^s \longrightarrow N$ through the quotient.

Moreover, if $\{g_1, \ldots, g_t\}$ is a generator system of B as a left R-module, with $g_k = (g_{k1}, .., g_{ks})$, then

$$\{\ (e'_j g_{k1}, \ldots, e'_j g_{ks})\ \}_{\substack{1 \le k \le t \\ 1 \le j \le m}} \ \cup\ \{\ (0_{R^m}, \ldots, \overset{i}{h_l}, \ldots, 0_{R^m})\ \}_{\substack{1 \le i \le s, \\ 1 \le l \le r}} \tag{4}$$

is a generator system of $\alpha^{-1}(T) = \alpha^{-1}(R^m \otimes_R B) + L^s$ as a left R-module, since

$$\{\ e'_j \otimes g_k\ \}_{\substack{1 \le j \le m, \\ 1 \le k \le t}} \ \cup\ \{\ h_l \otimes e_i\ \}_{\substack{1 \le l \le r, \\ 1 \le i \le s}}$$

is a generator system of $T = R^m \otimes_R B + L \otimes_R R^s$ and

$$\begin{array}{rcl} \alpha^{-1}(T) & = & \alpha^{-1}(R^m \otimes_R B) + \alpha^{-1}(L \otimes_R R^s) \\ & = & \alpha^{-1}(R^m \otimes_R B) + L^s \\ & = & {}_R\langle\ \alpha^{-1}(e'_j \otimes g_k),\ \alpha^{-1}(h_l \otimes e_i)\ \rangle_{i,j,k,l} \\ & = & {}_R\langle\ (g_{k1}e'_j, \ldots, g_{ks}e'_j), (0_{R^m}, \ldots, \overset{i}{h_l}, \ldots, 0_{R^m})\ \rangle_{i,j,k,l}. \end{array}$$

2 Computing $\mathrm{Tor}_k(M, N)$

Now, let $R = k\{x_1, \ldots, x_n;\ Q, Q', \preceq\}$ be a left PBW ring and N a finitely generated left R-module. Let

$$\cdots \xrightarrow{\partial_{k+1}} R^{s_k} \xrightarrow{\partial_k} R^{s_{k-1}} \longrightarrow \cdots \xrightarrow{\partial_1} R^{s_0} \xrightarrow{\partial_0} N \longrightarrow 0 \tag{5}$$

be a free resolution of N, where ∂_k is the matrix

$$\begin{pmatrix} g^k_1 \\ \vdots \\ g^k_{s_k} \end{pmatrix}$$

with $g_i^k = (g_{i1}^k, \ldots, g_{is_{k-1}}^k) \in R^{s_{k-1}}$ for $k \geq 1$ and $1 \leq i \leq s_k$ (an algorithm to compute it may be found in [2, chapter 6]).

In this section we show a method for computing $\mathrm{Tor}_k(M, N)$ when M is a finitely presented R-bimodule, i.e., we show how to compute the k-th homology module of the complex

$$\cdots \overset{1_M \otimes \partial_{k+1}}{\longrightarrow} M \otimes_R R^{s_k} \overset{1_M \otimes \partial_k}{\longrightarrow} \cdots \overset{1_M \otimes \partial_1}{\longrightarrow} M \otimes_R R^{s_0} \longrightarrow 0. \tag{6}$$

Remark 2.1. The case $k = 0$ may be treated apart, since $\mathrm{Tor}_0(M, N) \cong M \otimes_R N$.

We start with the presentations $0 \to L \to R^m \to M \to 0$ and $0 \to \ker \partial_0 \to R^{s_0} \to N \to 0$ of the R-bimodule M, resp. of the left R-module N.

Since $\{g_1^1, \ldots, g_{s_1}^1\}$ is a generator system of $\ker \partial_0 = \mathrm{im}\, \partial_1$ as a left R-module and

$$\gamma : R^{ms_0}/(\alpha^{-1}(R^m \otimes_R \ker \partial_0) + L^{s_0}) \longrightarrow \mathrm{Tor}_0(M, N),$$

as in (3) is an isomorphism of left R-modules, we completely describe a presentation of $\mathrm{Tor}_0(M, N)$ by giving the generator system

$$\{\ (e'_j g_{l1}^1, \ldots, e'_j g_{ls_0}^1)\ \}_{\substack{1 \leq l \leq s_1,\\ 1 \leq j \leq m}} \ \cup\ \{\ (0_{R^m}, \ldots, \overset{i}{h_l}, \ldots, 0_{R^m})\ \}_{\substack{1 \leq i \leq s_0,\\ 1 \leq l \leq r}}$$

of $\alpha^{-1}(R^m \otimes_R \ker \partial_0) + L^{s_0}$, where $\{h_1, \ldots, h_r\}$ is a two-sided Gröbner basis of M (see (4)).

Let us return to the general case. Consider again the presentation $0 \to L \to R^m \overset{p_M}{\to} M \to 0$ of the R-bimodule M and the free resolution (5) of N.

For all $k \geq 1$, let $d_k = \gamma_{k-1}^{-1} \circ (1_M \otimes \partial_k) \circ \gamma_k$, where $\gamma_k : R^{ms_k}/L^{s_k} \longrightarrow M \otimes_R R^{s_k}$ is the isomorphism defined as in (3) with $N = R^{s_k}$.

Clearly, the complex

$$\cdots \longrightarrow R^{ms_k}/L^{s_k} \xrightarrow{d_k} R^{ms_{k-1}}/L^{s_{k-1}} \xrightarrow{d_{k-1}} \cdots \xrightarrow{d_1} R^{ms_0}/L^{s_0} \longrightarrow 0 \tag{7}$$

is isomorphic to the one in (6), so $\mathrm{Tor}_k(M, N)$ may be computed as the homology of (7).

By definition, for all $1 \leq i \leq s_k$, $1 \leq j \leq m$,

$$\begin{aligned} d_k(e_{ij} + L^{s_k}) &= \gamma_{k-1}^{-1}(1_M \otimes \partial_k)\, \gamma_k(e_{ij} + L^{s_k}) \\ &= \gamma_{k-1}^{-1}((1_M \otimes \partial_k)\, (p_M(e'_j) \otimes e_i)) \\ &= \gamma_{k-1}^{-1}(p_M(e'_j) \otimes (g_{i1}^k, \ldots, g_{is_{k-1}}^k)) \\ &= (g_{i1}^k e'_j, \ldots, g_{is_{k-1}}^k e'_j) + L^{s_{k-1}}. \end{aligned} \tag{8}$$

Let $\widetilde{d_k} : R^{ms_k} \longrightarrow R^{ms_{k-1}}$ be the block-built matrix

$$A_k = \left(\begin{array}{c|c|c} g_{11}^k I_m & \cdots & g_{1s_{k-1}}^k I_m \\ \hline \vdots & & \vdots \\ \hline g_{s_k 1}^k I_m & \cdots & g_{s_k s_{k-1}}^k I_m \end{array} \right) \in \mathcal{M}_{ms_k \times ms_{k-1}}(R),$$

where I_m denotes the $m \times m$-identity matrix. Since A_k is built of blocks which are elements of R times the identity matrix, we have $\widetilde{d_k}(L^{s_k}) \subseteq L^{s_{k-1}}$, and the diagram

$$\begin{tikzcd} R^{ms_k} \arrow[r, "\pi_k"] \arrow[d, "\widetilde{d_k}"'] & R^{ms_k}/L^{s_k} \arrow[d, "d_k"] \\ R^{ms_{k-1}} \arrow[r, "\pi_{k-1}"'] & R^{ms_{k-1}}/L^{s_{k-1}} \end{tikzcd}$$

is commutative.

The above discussion proves the following result:

Theorem 2.2. *With the previous notation, for all* $k \geq 1$

1. $\ker d_k = \ker \pi_{k-1}\widetilde{d_k}/L^{s_k}$*;*
2. $\operatorname{im} d_k = \operatorname{im} \pi_{k-1}\widetilde{d_k} \subseteq R^{ms_{k-1}}/L^{s_{k-1}}$ *is generated by*

   $$\{(g_{i1}^k e'_j, \ldots, g_{is_{k-1}}^k e'_j) + L^{s_{k-1}}\}_{\substack{1\leq i \leq s_k \\ 1 \leq j \leq m}}$$

   as a left R-module (note that $(g_{i1}^k e'_j, \ldots, g_{is_{k-1}}^k e'_j)$ *is the* $(j+m(i-1))$*-th row of* A_k*);*
3. $\operatorname{Tor}_k(M,N) \cong \ker d_k / \operatorname{im} d_{k+1} = \ker \pi_{k-1}\widetilde{d_k}/({}_R\langle \text{rows of } A_{k+1}\rangle + L^{s_k})$.

Remark 2.3. To compute $\operatorname{Tor}_k(M,N)$ we start with a finite presentation $0 \to L \to R^m \overset{p_M}{\to} M \to 0$ of the R-bimodule M (say, e.g., we have computed a two-sided Gröbner basis $\{h_1, \ldots, h_r\} \subseteq R^m$ of the R bimodule L) and a free resolution of the left R-module N as (5).

The matrix A_k is block-built as above, using the matrix

$$\partial_k = \begin{pmatrix} g_1^k \\ \vdots \\ g_{s_k}^k \end{pmatrix}.$$

The set $\{\ (0_{R^m}, \ldots, \overset{i}{h_l}, \ldots, 0_{R^m})\ \}_{i,l=1}^{s_k,r}$ is a generator system (in fact, it is a two-sided Gröbner basis when $\{h_1, \ldots, h_r\}$ so is) of L^{s_k} as an R-bimodule.

Then, we compute the kernel of $\pi_{k-1}\widetilde{d_k} : R^{ms_k} \longrightarrow R^{ms_{k-1}}/L^{s_{k-1}}$ using syzygies. Indeed (see [1]), let H_k be the matrix

$$H_k = \left(\frac{A_k}{A'_k} \right) \in \mathcal{M}_{(ms_k + rs_{k-1}) \times ms_{k-1}}(R),$$

where A'_k is the matrix whose rows are the generators of $L^{s_{k-1}}$ as a left R-module. Then, if

$$Syz^l(H_k) = \ _R\langle p_1, ..., p_l \rangle, \quad \text{with } p_i = (p'_i, p''_i) \in R^{ms_k} \times R^{rs_{k-1}},$$

then $\ker \pi_{k-1}\widetilde{d_k} \ = \ _R\langle p'_1, ..., p'_l \rangle$.

Example 2.4. *Let $R = U(\mathfrak{sl}(2))$, the universal enveloping algebra of the Lie algebra of traceless 2×2-matrices, where $k = \mathbb{C}$ (or $k = \mathbb{Q}$). We know (see, e.g., [2]) that R is the PBW algebra $k\{x, y, z;\ Q\ \preceq_\omega\}$ with $Q = \{\ yx - xy + z,\ zx - xz - 2x,\ zy - yz + 2y\ \}$ and, say, $\omega = (1, 2, 2)$.*

Let $N = R^2/B$, where B is the left R-module generated by

$$g_1 = (y^3, x),\ g_2 = (y, xz),\ g_3 = (0, xy^2z - 2yz^2 + 2yz - x).$$

The left syzygy module $Syz^l(g_1, g_2, g_3)$ is generated by $g = (1, -y^2, 1) \in R^3$, and hence

$$0 \longrightarrow R \xrightarrow{\delta_1} R^3 \xrightarrow{\delta_0} N \longrightarrow 0, \tag{9}$$

where $\partial_0 = \begin{pmatrix} g_1 \\ g_2 \\ g_3 \end{pmatrix}$ and $\partial_1 = (g)$, is a free resolution of N.

Let L be the R-bimodule generated by $\{(C, 1), (1, C)\}$, where C is the Casimir element $z^2/2 + 2xy - z$ (a well known central element of $U(\mathfrak{sl}(2))$), and let $M = R^2/L$.

For all $k \geq 2$ we have $\mathrm{Tor}_k(M, N) = 0$, as the free resolution (9) of N has length 2.

For $k = 0$, we have (see 2.1) $\mathrm{Tor}_0(M, N) \cong R^6\ /\ (\alpha^{-1}(R^2 \otimes_R B) + L^3)$, and

$$\{\ (1, 0, -y^2, 0, 1, 0)\ ,\ (0, 1, 0, -y^2, 0, 1)\ ,\ (C, 1, 0, 0, 0, 0)\ ,\ (0, 0, C, 1, 0, 0)\ ,$$
$$(0, 0, 0, 0, C, 1)\ ,\ (1, C, 0, 0, 0, 0)\ ,\ (0, 0, 1, C, 0, 0)\ ,\ (0, 0, 0, 0, 1, C)\ \}$$

is a generator system of $\alpha^{-1}(R^2 \otimes_R B) + L^3$ as a left R-module.

For $k = 1$, we have $\mathrm{Tor}_1(M, N) \cong \ker(\pi_0\widetilde{d_1})\ /\ ({}_R\langle \text{rows of } A_2 \rangle + L)$, but in this particular example $A_2 = 0$ since $\partial_2 = 0$.

As pointed out in 2.3, $\ker \pi_0 \widetilde{d_1}$ *is obtained from the left syzygy module of the rows of*

$$A_1 = \begin{pmatrix} 1 & 0 & -y^2 & 0 & 1 & 0 \\ 0 & 1 & 0 & -y^2 & 0 & 1 \end{pmatrix}$$

and the generators of L^3. *Indeed, by picking up just the first two components of each element of the generator system*

$$\begin{aligned} &\{(z^2+4xy-2z, 2, -2, 2y^2, -2, 0, 0, 0)\ , \\ &\ (1, z^2/2+2xy-z, 0, 0, 0, -1, y^2, -1)\ , \\ &\ (8xy-4z, -4z^4-4xyz^2+2z^3+4, -4, 4y^2, -4, 2y^2, \\ &\qquad\qquad -2y^2z^2+16y^2z-32y^2, 2z^2)\} \end{aligned}$$

of the left syzygy module, we obtain the generator system

$$\{\ (2C, 2)\ ,\ (1, C)\ ,\ (8xy-4z, -z^4-4xyz^2+2z^3+4)\ \}$$

of $\ker \pi_0 \widetilde{d_1}$. *Therefore,* $\mathrm{Tor}_1(M, N) \cong \ker \pi_0 \widetilde{d_1}/L = 0$, *since* $(8xy-4z, -z^4-4xyz^2+2z^3+4) \in L$ *(one may check this out by dividing the element by a two-sided Gröbner basis of* L*).*

Acknowledgement

Both authors would like to thank Pepe Bueso for his useful suggestions.

References

[1] Bueso, J.L.; Castro, F.; Gómez Torrecillas, J.; Lobillo, F.J. Homological computations in PBW modules. Alg. Rep. Theory **2001**, 4 (2), 201-218.

[2] Bueso, J.L.; Gomez Torrecillas, J.; Verschoren, A. Algorithmic Methods in Noncommutative Algebra: Applications to Quantum Groups; Kluwer: Dordrecht, The Netherlands, 2003.

[3] García Román, S.; García Román, M. Gröbner bases and syzygies on bimodules, preprint. `math.RA/0405550`

[4] Smith, G. Computing global extension modules. J. Symb. Comput. **2000**, 29 (4-5), 729-746.